

Reference List

1. Craig JW, Glick C, Phillips R, Hall SL, Smith J, Browne J. Recommendations for involving the family in developmental care of the NICU baby. *Journal of perinatology*. 2015;35:S5-S8.
2. Feldman R, Eidelman AI, Sirota L, Weller A. Comparison of skin-to-skin (kangaroo) and traditional care: parenting outcomes and preterm infant development. *Pediatrics*. 2002;110(1 Pt 1):16-26.
3. Lester BM, Salisbury AL, Hawes K, Dansereau LM, Bigsby R, Laptook A, et al. 18-Month follow-up of infants cared for in a single-family room Neonatal Intensive Care Unit. *J Pediatr*. 2016;177:84-9.
4. Welch MG, Meeka S, Halperin MS, Austin J, Stark RI, Hofer MA, et al. Depression and anxiety symptoms of mothers of preterm infants are decreased at 4 months corrected age with family nurture intervention in the NICU. *Arch womens mental health*. 2016;19(1):51-61.
5. Cheng C, Franck LS, Ye XY, Hutingson SA, Lee SK, O'Brien K, et al. Evaluating the effect of family integrated care on maternal stress and anxiety in Neonatal Intensive Care Units. *Journal of reproductive and infant psychology*. 2019;10:1-14.
6. Aftyka A, Rybojad B, Rosa W, Wrobel A, Karakula-Juchnowicz H. Risk factors for the development of posttraumatic stress disorder and coping strategies in mothers and fathers following infant hospitalization in the neonatal intensive care unit. *Journal of clinical nursing*. 2017;26:338-47.

7. Forcada-Guex M, Borghini A, Pierrehumbert B, Ansermet F, Muller-Nix C. Prematurity, maternal posttraumatic stress and consequences on the mother-infant relationship. *Early hum dev.* 2011;87(1):21-6.
8. Gondwe KW, Brandon D, Yang Q, Chirwa E, Holditch-Davis D. Validation of the Chichewa perinatal PTSD questionnaire and Chichewa child health worry scale. *International journal of Africa nursing sciences.* 2018;9.:42-8.
9. Holditch-Davis D, Miles MS, Weaver MA, Black B, Beeber L, Hoyre S, et al. Patterns of distress in African-American mothers of preterm infants. *Journal of developmental and behavioral pediatric.* 2009;30(3):193-205.
10. Kim WJ, Lee E, Kim KR, Namkoong K, Park ES, Rha D. Progress of PTSD symptoms following birth: a prospective study in mothers of high-risk infants. *J perinatology.* 2015;35(8):575-9.
11. Hall SL, Phillips R, Hynan M. Transforming NICU care to provide comprehensive family support. *Newborn and infant nursing reviews.* 2016;16:69-73.
12. Hall SL, Hynan MT, Phillips R, Lassen S, Craig JW, Gover E, et al. The neonatal intensive parenting unit: an introduction. *J perinatol.* 2017;37:1259-64.
13. Hynan MT, Hall SL. Psychosocial program standards for NICU parents. *J perinatol.* 2015;35 Suppl 1:S1-4.
14. Hynan MT, Steinberg Z, Baker L, Cicco R, Geller PA, Lassen S, et al. Recommendations for mental health professionals in the NICU. *J perinatol.* 2015;35 Suppl 1:S14-8.
15. Hoffman MC, Mazzoni SE, Wagner BD, Laudenslager ML, Ross RG. Measures of maternal stress and mood in relation to preterm birth. *Obstet Gynecol.* 2016;3(3):545-52.

16. Holditch-Davis D, Santos H, Levy J, White-Traut R, O'Shea M, V. G, et al. Patterns of psychological distress in mothers of preterm infants. *Infant behavior and development*. 2015;41:154-63.
17. Hynan M, Mounts K, Vanderbilt D. Screening parents of high-risk infants for emotional distress: rationale and recommendations. *J perinatol*. 2013;33(748-53):748-53.
18. Bonanno GA, Westphal M, Anthony D, Mancini AD. Resilience to loss and potential trauma. *Annual review of clinical psychology*. 2011;7:511-35.
19. Kazak AE. Pediatric psychosocial preventative health model (ppphm): Research, practice, and collaboration in pediatric family systems medicine. *Family system health*. 2006;24:381-95.
20. Hall S, Ryan D, Beatty J, Grubbs L. Recommendations for peer-to-peer support for NICU parents. *Journal of perinatology*. 2015;35:S9-S13.
21. Melnyk B, Feinstein N, Alpert-Gillis L, Fairbanks E, Crean H, Sinkin RA, et al. Reducing premature infants' length of stay and improving parents' mental health outcomes with the Creating Opportunities for Parent Empowerment (COPE) neonatal intensive care unit program: a randomized, controlled trial. *Pediatrics*. 2006;118(5):e1414–e27.
22. Preyde M, Ardal F. Effectiveness of a parent 'buddy' program for mothers of very preterm infants in a neonatal intensive care unit. *CMAJ*. 2003;168:969-73.
23. Segre L, O'Hara M, Arndt S, Tatano-Beck C. Nursing care for postpartum depression, part 1: do nurses think they should offer both screening and counseling? *Am J matern child nurs*. 2010;35(4):220-5.

24. Segre L, Chuffo-Siewert R, Brock R, O'Hara M. Emotional distress in mothers of preterm hospitalized infants: a feasibility trial of nurse-delivered treatment. *J perinatol*. 2013;33(12):924-8.
25. Jotzo M, Poets C. Helping parents cope with the trauma of premature birth: an evaluation of a trauma-preventive psychological intervention. *Pediatrics*. 2005;115(4):915-19.
26. Shaw R, St. John N, E. L, Jo B, Benitez W, Stevenson DK, et al. Prevention of traumatic stress in mothers of preterm infants: a randomized controlled trial. *Pediatrics*. 2013;132(4):1-9.
27. Shaw R, St. John N, Lilo E, Jo B, Benitez W, Stevenson D, et al. Prevention of traumatic stress in mothers of preterms: 6-month outcomes. *Pediatrics*. 2014;134:e481-88.
28. Anand KJ. Clinical importance of pain and stress in preterm neonates. *Biology of the neonate*. 1998;73(1):1-9.
29. Anand KJ, Aranda JV, Berde CB, Buckman S, Capparelli EV, Carlo W, et al. Summary proceedings from the Neonatal Pain-Control Group. *Pediatrics*. 2006;117(3 Pt 2):S9-S22.
30. Carbajal R, Rousset A, Danan C, Coquery S, Nolent P, Ducrocq S, et al. Epidemiology and treatment of painful procedures in neonates in intensive care units. *Jama*. 2008;300(1):60-70.
31. Vinall J, Grunau RE. Impact of repeated procedural pain-related stress in infants born very preterm. *Pediatr res*. 2014;75(5):584-7.
32. Doesburg SM, Chau CM, Cheung TP, Moiseev A, Ribary U, Herdman AT, et al. Neonatal pain-related stress, functional cortical activity and visual-perceptual abilities in school-age children born at extremely low gestational age. *Pain*. 2013;154(10):1946-52.
33. Hermann C, Hohmeister J, Demirakca S, Zohsel K, Flor H. Long-term alteration of pain sensitivity in school-aged children with early pain experiences. *Pain*. 2006;125(3):278-85.

34. Holsti L, Grunau RE, Oberlander TF, Whitfield MF. Prior pain induces heightened motor responses during clustered care in preterm infants in the NICU. *Early hum dev.* 2005;81(3):293-302.
35. Simons SH, van Dijk M, Anand KS, Roofthoof D, van Lingen RA, Tibboel D. Do we still hurt newborn babies? A prospective study of procedural pain and analgesia in neonates. *Archives of pediatrics & adolescent medicine.* 2003;157(11):1058-64.
36. Morrow C, Hidinger A, Wilkinson-Faulk D. Reducing neonatal pain during routine heel lance procedures. *Am J matern child nurs.* 2010;35(6):346-54; quiz 54-6.
37. Obeidat H, Kahalaf I, Callister LC, Froelicher ES. Use of facilitated tucking for nonpharmacological pain management in preterm infants: a systematic review. *J perinat neonatal nurs.* 2009;23(4):372-7.
38. Liaw JJ, Yang L, Katherine Wang KW, Chen CM, Chang YC, Yin T. Non-nutritive sucking and facilitated tucking relieve preterm infant pain during heel-stick procedures: a prospective, randomised controlled crossover trial. *Int J nurs stud.* 2012;49(3):300-9.
39. Pillai Riddell R, Racine NM, Turcotte K, Uman L, Horton R, Din Osmun L, et al. Nonpharmacological management of procedural pain in infants and young children: an abridged Cochrane review. *Pain res manag.* 2011;16(5):321-30.
40. Axelin A, Anderzen-Carlsson A, Eriksson M, Polkki T, Korhonen A, Franck LS. Neonatal Intensive Care nurses' perceptions of parental participation in infant pain management: a comparative focus group study. *J perinat neonatal nurs.* 2015;29(4):363-74.

41. Milgrom J, Newnham C, Anderson P, Doyle P, Gemmill A, Lee K, et al. Early sensitivity training for parents of preterm infants: impact on the developing brain. *Pediatr res*. 2010;67(3):330-5.
42. Ortenstrand A, Westrup B, Brostrom EB, Sarman I, Akerstrom S, Brune T, et al. The Stockholm Neonatal Family Centered Care Study: effects on length of stay and infant morbidity. *Pediatrics*. 2010;125(2):e278-85.
43. Shah PS, Herbozo C, Aliwalas LL, Shah VS. Breastfeeding or breast milk for procedural pain in neonates. *Cochrane database syst rev*. 2012;12:Cd004950.
44. Johnston C, Campbell-Yeo M, Disher T, Benoit B, Fernandes A, Streiner D, et al. Skin-to-skin care for procedural pain in neonates. *Cochrane database syst rev*. 2017;2:Cd008435.
45. Yin T, Yang L, Lee TY, Li CC, Hua YM, Liaw JJ. Development of atraumatic heel-stick procedures by combined treatment with non-nutritive sucking, oral sucrose, and facilitated tucking: a randomized controlled trial. *Int J nurs stud*. 2015;52(8):1288-9.
46. American Academy of Pediatrics (AAP) Task Force on Circumcision. Male circumcision. *Pediatrics*. 2012;130(3):e756-85.
47. Kumar P, Denson SE, Mancuso TJ. Premedication for nonemergency endotracheal intubation in the neonate. *Pediatrics*. 2010;125(3):608-15.
48. American Academy of Pediatrics (AAP) Committee on Fetus and Newborn and Section on Anesthesiology and Pain Management. Prevention and management of procedural pain in the neonate: an update. *Pediatrics*. 2016;137(2):1-13.
49. Bellu R, de Waal KA, Zanini R. Opioids for neonates receiving mechanical ventilation. *Cochrane database of systematic reviews*. 2008.

50. Anand KJ, Barton BA, McIntosh N, Lagercrantz H, Pelausa E, Young TE, et al. Analgesia and sedation in preterm neonates who require ventilatory support: results from the NOPAIN trial. *Neonatal Outcome and Prolonged Analgesia in Neonates. Archives of pediatrics & adolescent medicine.* 1999;153(4):331-8.
51. Aranda JV, Carlo W, Hummel P, Thomas R, Lehr VT, Anand KJ. Analgesia and sedation during mechanical ventilation in neonates. *Clinical therapeutics.* 2005;27(6):877-99.
52. Ng E, Taddio A, Ohlsson A. Intravenous midazolam infusion for sedation of infants in the neonatal intensive care unit. *Cochrane database syst rev.* 2017;1:Cd002052.
53. Laudert S, Liu WF, Blackington S, Perkins B, Martin S, Macmillan-York E, et al. Implementing potentially better practices to support the neurodevelopment of infants in the NICU. *J perinatol.* 2007;27 Suppl 2:S75-93.
54. Milgrom J, Newnham C, Martin PR, Anderson PJ, Doyle LW, Hunt RW. Early communication in preterm infants following intervention in the NICU. *Early human dev.* 2013;89(9):755-67.